

# The University of Bradford Institutional Repository

<http://bradscholars.brad.ac.uk>

This work is made available online in accordance with publisher policies. Please refer to the repository record for this item and our Policy Document available from the repository home page for further information.

To see the final version of this work please visit the publisher's website. Available access to the published online version may require a subscription.

**Link to Publisher's version:** <http://dx.doi.org/10.1167/iovs.14-15308>

**Citation:** Deshpande P, Ortega I, Sefat F, Sangwan V S, Green N, Claeysens F and MacNeil S (2015) Rocking Media Over Ex Vivo Corneas Improves This Model and Allows the Study of the Effect of Proinflammatory Cytokines on Wound Healing. *Investigative Ophthalmology & Visual Science*. 56(3): 1553-61.

**Copyright statement:** © 2015 The Authors. Published by ARVO. This work is licensed under a Creative Commons Attribution-NonCommercial-NoDerivatives 4.0 International License. (<http://creativecommons.org/licenses/by-nc-nd/4.0/>).



# Rocking Media Over Ex Vivo Corneas Improves This Model and Allows the Study of the Effect of Proinflammatory Cytokines on Wound Healing

Pallavi Deshpande,<sup>1</sup> Ílida Ortega,<sup>1</sup> Farshid Sefat,<sup>1</sup> Virender S. Sangwan,<sup>2</sup> Nicola Green,<sup>1</sup> Frederik Claeysens,<sup>1</sup> and Sheila MacNeil<sup>1</sup>

<sup>1</sup>Department of Materials Science and Engineering, Kroto Research Institute, University of Sheffield, Sheffield, United Kingdom  
<sup>2</sup>Sudhakar and Sreekanth Ravi Stem Cell Laboratory, LV Prasad Eye Institute, Kallam Anji Reddy Campus, LV Prasad Marg, Hyderabad, Telangana, India

Correspondence: Sheila MacNeil, Department of Materials Science and Engineering, Kroto Research Institute, North Campus, University of Sheffield, Broad Lane, Sheffield S3 7HQ, UK; s.macneil@sheffield.ac.uk.

Submitted: July 23, 2014  
 Accepted: December 8, 2014

Citation: Deshpande P, Ortega Í, Sefat F, et al. Rocking media over ex vivo corneas improves this model and allows the study of the effect of proinflammatory cytokines on wound healing. *Invest Ophthalmol Vis Sci.* 2015;56:1553–1561. DOI:10.1167/ iovs.14-15308

**PURPOSE.** The aim of this work was to develop an in vitro cornea model to study the effect of proinflammatory cytokines on wound healing.

**METHODS.** Initial studies investigated how to maintain the ex vivo models for up to 4 weeks without loss of epithelium. To study the effect of cytokines, corneas were cultured with the interleukins IL-17A, IL-22, or a combination of IL-17A and IL-22, or lipopolysaccharide (LPS). The effect of IL-17A on wound healing was then examined.

**RESULTS.** With static culture conditions, organ cultures deteriorated within 2 weeks. With gentle rocking of media over the corneas and carbon dioxide perfusion, the ex vivo models survived for up to 4 weeks without loss of epithelium. The cytokine that caused the most damage to the cornea was IL-17A. Under static conditions, wound healing of the central corneal epithelium occurred within 9 days, but only a single-layered epithelium formed whether the cornea was exposed to IL-17A or not. With rocking of media gently over the corneas, a multilayered epithelium was achieved 9 days after wounding. In the presence of IL-17A, however, there was no wound healing evident. Characterization of the cells showed that wherever epithelium was present, both differentiated cells and highly proliferative cells were present.

**CONCLUSIONS.** We propose that introducing rocking to extend the effective working life of this model and the introduction of IL-17A to this model to induce aspects of inflammation extend its usefulness to study the effects of agents that influence corneal regeneration under normal and inflamed conditions.

**Keywords:** cornea organ culture, inflammation, wound healing

The cornea, through which light enters the eye, is a transparent, avascular tissue<sup>1</sup> that is responsible for keeping bacteria from entering into the eye<sup>2</sup> while at the same time allowing oxygen to enter.<sup>3</sup> This is achieved through tight junctions present at the superficial layer of the cornea, creating a barrier.<sup>4</sup> If violated, this barrier function makes the cornea and possibly the whole eye susceptible to infection.<sup>5</sup>


Physical damage to the corneal epithelium is the most common cause of wounding. Wound healing of the cornea involves the removal of necrotic cells, infiltration of neutrophils, and migration of cells from the wound edge covering the wound, along with simultaneous multilayer formation.<sup>6</sup>

Corneal injuries are usually associated with some level of inflammation that includes neutrophils,<sup>7</sup> macrophages, dendritic cells, and lymphocytes.<sup>8</sup> Neutrophils are recruited to the injury site from the limbal blood vessels<sup>7</sup> and tear fluid<sup>6</sup> just after stromal apoptosis takes place<sup>9</sup> due to the release of IL-1<sup>10</sup> and TNF- $\alpha$ <sup>11</sup> by the injured epithelial cells. The neutrophils phagocytose cellular debris and pathogens,<sup>12,13</sup> clearing up the site of injury, and also release large amounts of cytotoxic agents that assist in the phagocytosis of microbes.<sup>14</sup>

Most cornea organ culture models employed to study the transfer of cells to the cornea or the effects of chemicals on the corneal epithelium and wound healing<sup>15–19</sup> have a life span of around 3 to 4 weeks at maximum, and most of these models are static cultures.<sup>15,16,20</sup>

The static models, while fairly simple to set up, do not mimic the situation in vivo in which the corneas are kept intermittently moist, through the blinking action of the eyelids, but are not submerged. In the current study the corneas were placed in a simple rocking system (based on a commercially available egg incubator modified for CO<sub>2</sub> perfusion). Using this technique, multiple corneas could be cultured simultaneously, and we demonstrate that the models survive for at least 4 weeks, which is twice as long as under static conditions.<sup>15</sup>

We then used these conditions to develop an in vitro cornea wound model and incorporated cytokines known to induce inflammation. We identified the cytokine IL-17A, which induced clear tissue disruption to the corneal epithelium, and studied wound healing in the model in the presence of this cytokine over 9 days under both static and rocking culture conditions. We show that rocking provides a model in which



**FIGURE 1.** Photograph of a commercially available egg incubator on a commercially available platform rocker (A) and schematic of the cornea organ culture in a Petri dish at a (i) horizontal position and (ii) inclined position on a platform rocker (B).

wound healing occurs rapidly except when in the presence of IL-17A.

## MATERIALS AND METHODS

### Cornea Organ Culture

New Zealand white rabbits were obtained from a rabbit farm (Hook Farm, Hampshire, UK). The rabbits weighed between 2.4 and 2.6 kg. Excess fat and tissue were excised from the eyes, which were then processed as described previously.<sup>16</sup> The corneoscleral buttons were cultured at an air-liquid interface in 1 mL medium consisting of Dulbecco's modified Eagle's medium (DMEM GlutaMAX; Gibco Life Technologies Ltd., Paisley, UK) and Ham's F12 medium (Biosera, Ringmer, UK) in a 1:1 ratio supplemented with 10% fetal calf serum (Biosera), 10 ng/mL epidermal growth factor (EGF), 5 µg/mL insulin, 2.5 µg/mL Amphotericin B, 100 IU/mL penicillin, and 100 µg/mL streptomycin (Sigma Aldrich, Poole, UK). The corneas were cultured at 37°C in a humidified, carbon dioxide (CO<sub>2</sub>) environment.

### Static Culture of Corneas

The medium was changed every other day, and the corneas were kept wet by dropping medium on the surface every day.

### Application of Cytokines to the Cornea

On day 7, the medium of the unwounded corneas was changed to contain 20 ng/mL recombinant human IL-17A (Peprotech, London, UK), 25 ng/mL IL-22 (Peprotech), a combination of IL-17A and IL-22 (20 ng/mL and 25 ng/mL, respectively), or 1 µg/mL lipopolysaccharide (LPS). All cytokines were added to media, and control organ cultures were cultured in media without cytokines. The medium was changed every other day with wetting of the corneas every day. On day 14, the samples

were fixed using 3.7% buffered formaldehyde and processed for conventional histology.

### Rocking Culture of the Corneas

After preparation of the corneoscleral button, corneas were placed at the periphery of a 90-mm Petri dish and held in place by the end of a pipette tip glued onto the Petri dish using Araldite epoxy adhesive (Araldite; Huntsman Advanced Materials, Basel, Switzerland). The corneas were cultured in 13 mL culture medium and placed in our in-house rocking system consisting of an egg incubator (R-COM King Suro Egg Incubator; P&T Poultry, Powys, UK), which was subsequently placed on a platform rocker (Platform Rocker STR6; Stuart, Stone, UK) (Fig. 1) set at a speed of 10 rocks per minute and maximum tilt angle of 6°. The corneas were maintained at 37°C in the humidified chamber of the egg incubator (Fig. 1). Sixteen corneas could be handled in the incubator at one time.

### Assessment of the Effect of Including a CO<sub>2</sub> Supply in the In-House Rocking Incubator

Cornea organ cultures were cultured in the in-house rocking incubator in the absence and presence of 5% CO<sub>2</sub>. CO<sub>2</sub> was added through a hole at the top of the egg incubator via tubing from a cylinder containing 5% CO<sub>2</sub> (see Fig. 1). The corneas were cultured for either 2 or 4 weeks and compared to corneas cultured under static conditions in a conventional cell incubator (in the presence of CO<sub>2</sub>) with the media changed every week.

### Corneal Epithelial Wounding

After cleaning the eyes of excess tissue, the corneas were held in position and the central area of the cornea was wounded using 20% methanol within an 8-mm-diameter trephine. After 40 seconds, the area was rinsed with PBS to remove any excess methanol; and using a sclerotome knife, the epithelium was




FIGURE 2. Schematic of an inflamed in vitro cornea wound model.

scraped from the treated area (Fig. 2). The corneoscleral buttons were excised and then processed to be placed in 90-mm Petri dishes as described previously in this paper. Organ cultures with unwounded epithelia were also set up as controls.

On day 2, the media of the wounded and unwounded corneas were changed to contain the cytokine that had the greatest effect on corneal morphology—IL-17A (Fig. 2). Controls were both wounded and unwounded corneas in organ culture medium without IL-17A. On day 7 the corneas in the rocker were transferred to 35-mm Petri dishes. On day 9 the corneas were fixed for 1 hour using 3.7% buffered formaldehyde and processed for conventional histology. This rocking protocol was compared to a previous protocol<sup>16</sup> in which corneas were cultured in a 35-mm Petri dish under static conditions in an incubator maintained at 37°C in a humidified, 5% CO<sub>2</sub> environment.

### Immunohistochemistry

Corneal sections of 4 μm were mounted onto Superfrost Plus microscopic slides (Thermo Fisher Scientific, Loughborough, UK) and stained for cytokeratin 3 (CK3) and P63 (stem cell marker) as described previously.<sup>16</sup> The samples were imaged using a Zeiss LSM 510 META confocal microscope (Carl Zeiss MicroImaging, Jena, Germany) at the Kroto Research Institute Confocal imaging facility.

## RESULTS

### Investigation of the Effect of Culturing Corneas Under Rocking Versus Static Conditions

We assessed the integrity of the corneal epithelium under static and rocking conditions. In order to mimic the blinking action of the eyelids and consequently keep the corneas wet, the corneas were placed on a rocker within an incubator maintained at 37°C with a small volume of media that flowed intermittently over the cornea as a result of the rocking action. These epithelia were compared to those of corneas cultured within static incubators at 37°C.

Figure 3 shows the effect of CO<sub>2</sub> on the corneas cultured under rocking conditions. It could be clearly seen that culturing the corneas under static conditions in the presence of CO<sub>2</sub> for 2 weeks (Fig. 3A) and 4 weeks (Fig.

3D) resulted in a epithelium one or two cell layers thick, while rocking conditions without (Fig. 3B) and with (Fig. 3C) CO<sub>2</sub> resulted in a epithelium five cell layers thick over the same period. On culturing the corneas under rocking conditions for 4 weeks, the epithelium was completely lost in the absence of CO<sub>2</sub> (Fig. 3E) but was retained in its presence (Fig. 3F). After 2 weeks it was observed that the organization of the epithelium was poor in the absence of CO<sub>2</sub> (Fig. 3B) compared to the corneas cultured in the presence of CO<sub>2</sub> (Fig. 3C); and after 4 weeks, even in the presence of CO<sub>2</sub> (Fig. 3F), the epithelium appeared to have lost the superficial desquamated cells compared to 2 weeks in culture (Fig. 3C).

### Effect of Inflammatory Cytokines on Corneal Epithelium

Previous studies<sup>21</sup> and unpublished studies in the group have shown that exposure to the cytokines IL-17A and IL-22 leads to inflammation of the epidermis, while a combination of IL-17A and IL-22 further enhanced inflammation of human colonic subepithelial myofibroblasts.<sup>22</sup> Similarly, LPS has been shown to induce inflammation of the epithelium.<sup>23,24</sup> Accordingly, these cytokines and LPS were selected as potential proinflammatory cytokines for this study.

The corneas were cultured for 7 days prior to the application of the cytokines. Figure 4 shows that IL-17A (Fig. 4B) or a mixture of IL-17A and IL-22 (Fig. 4D) resulted in the formation of a poor epithelium with a very loosely attached multilayer of cells. Addition of IL-17A alone led to a split in the epithelial multilayer. The basal cells remained well adhered to the corneal surface with a suprabasal split of the cells above this. The control cornea that had not been exposed to the cytokines showed a normal multilayer of cells on the corneal surface (Fig. 4A), while corneas exposed to IL-22 (Fig. 4C) and LPS (Fig. 4E) formed an epithelium that was similar to that seen in the control.

### Effect of Cytokines on Wound Healing

Corneas were initially cultured for 2 days prior to the application of IL-17A. Figure 5 shows the effect of IL-17A on wound healing of the cornea. Figure 5A shows a normal cornea under static conditions. Here there was no multilayer formation, and after 9 days in culture there was just a single layer of cells on the corneal surface. Interleukin-17A (Fig. 5B)






FIGURE 3. Hematoxylin and eosin of cornea organ cultures at 2 weeks under static conditions (A) and rocking conditions without (B) and with (C) CO<sub>2</sub> and at 4 weeks under static conditions (D) and rocking conditions without (E) and with (F) CO<sub>2</sub>.

on the unwounded cornea again resulted in a single layer of cells on the surface, with the cells appearing abnormal with some level of perinuclear vacuolation. The wounded cornea (Fig. 5C) was approximately one layer thick, and a similar epithelium was seen when the wounded cornea was exposed to IL-17A (Fig. 5D). Although the images show the central area of the cornea where wounding was carried out, the epithelium shown in the images is representative of the whole corneal epithelium as seen by visual examination.

Under rocking conditions, however, the control unwounded cornea had a much improved epithelium compared to the cornea cultured under static conditions, with approximately seven layers in thickness (Fig. 5E). Interleukin-17A had an adverse effect on the unwounded cornea resulting in the loss of most of the epithelium, and only a very few cells remained present on the surface (Fig. 5F). Nine days after wounding the control cornea, a multilayered epithelium had formed of approximately five cell layers thick under rocking conditions





FIGURE 4. Hematoxylin and eosin of cornea organ culture exposed to cytokines. (A) The control cornea without cytokine exposure; (B) exposed to IL-17A (20 ng/mL); (C) exposed to IL-22 (25 ng/mL); (D) exposed to both IL-17A and -22 (20 ng/mL and 25 ng/mL, respectively); and (E) exposed to lipopolysaccharide (1 µg/mL).



**FIGURE 5.** Hematoxylin and eosin of in vitro cornea organ culture after 9 days under static (A–D) and rocking (E–H) conditions. (A, E) Unwounded corneas; (B, F) unwounded corneas exposed to IL-17A; (C, G) a wounded cornea; and (D, H) a wounded cornea exposed to IL-17A.

(Fig. 5G); however, on exposing the wounded cornea to IL-17A, the cornea lost all the epithelial cells and there was no regeneration of epithelium visible even under rocking conditions (Fig. 5H). Also in the wounded corneas it was noted that the stromal cells were not present in the area where wounding had taken place (Figs. 5C–H).

Immunohistochemistry showed that the cells on the corneal surface were not affected by the addition of IL-17A. Cytokeratin 3 was expressed despite being exposed to IL-17A under both static and rocking conditions (Figs. 6A–G). P63 continued to be expressed in all the corneas whether exposed to IL-17A or not (Figs. 6I–O). However, this was more prominent at the peripheral area of the cornea than the central area. P63 could also be visualized in the basal layer of cells in the central corneal region, where a multilayer of cells formed in the unwounded (Figs. 6M, 7A–C) and wounded (Figs. 6O, 7D–F) corneas under rocking conditions, but could not be seen where only a monolayer of cells had formed (Figs. 6I–L, 6N). In the wounded corneas exposed to IL-17A under rocking conditions (Figs. 6H, 6P), no cells were present on the corneal surface, as seen using hematoxylin and eosin (H&E) staining; hence it was not possible to examine CK3 or P63 expression (Fig. 5).

## DISCUSSION

From the current study we suggest that the epithelia of three-dimensional (3D) cornea models can be improved by simply rocking the corneas from side to side to mimic the blinking action of the eye. Blinking provides the cornea with fresh tears and nutrients to the corneal cells.<sup>25</sup> Richard et al.<sup>19</sup> in 1991 placed corneas on a standard laboratory rocking platform where the intermittent movement of fluid over the cornea was used to mimic blinking. They compared the rocking technique to corneas cultured under static submerged conditions. Results showed that culturing the corneas intermittently at an air–liquid interface reduced epithelial, stromal, and endothelial intercellular edema. However, placing the rocker in an incubator does not seem to have been taken up by others—possibly because this can make the incubator system susceptible to overheating. In 2006, a more complex perfusion model<sup>18</sup> was published in which media were delivered drop-wise onto the cornea to mimic the blinking action instead.

In the current study the corneas were placed in a simple rocking system (based on a commercially available egg incubator modified for CO<sub>2</sub> perfusion). Using this technique, it was readily demonstrated that the 3D models could be maintained in culture for twice as long as under static conditions.


When culturing corneal limbal stem cells for use in transplantation, some groups have cultured cells on an amniotic membrane at an air–liquid interface<sup>26</sup> to form a multilayer of cells prior to transplantation, while others have cultured the cells under submerged conditions<sup>27</sup> forming only a monolayer of cells, and both report clinical success. With the current technique it would be possible to place an amniotic membrane or a synthetic membrane on the cornea and look at the formation of an epithelium from cultured cells, or indeed from explants of tissue under rocking conditions, to see which provides best epithelial regeneration in vitro.

In order to establish the model, the effect of including a CO<sub>2</sub> supply in the rocking incubator was first studied. The presence of CO<sub>2</sub> was crucial for pH maintenance around 6.9 to 7.8, which is important for cells to grow.<sup>28</sup> In cells cultured in bicarbonate-buffered media, the bicarbonate leaves the medium in the form of carbon dioxide, making the medium very alkaline. The presence of 5% CO<sub>2</sub> in the incubator maintains an equilibrium in the tissue culture dish, preventing the increase in the pH to over 8.5,<sup>29</sup> which is toxic to the cells.<sup>30</sup> Our results, as expected, confirmed the necessity of a CO<sub>2</sub> supply to our in-house incubator, especially while culturing these corneas for long periods of time (4 weeks) compared to shorter periods (2 weeks).

In the immune system, T helper cell functions<sup>31</sup> can be divided into Th1, Th2, and Th17.<sup>32</sup> Th-17 is a T helper cell subset that on differentiation releases IL-17, IL-21, and IL-22 with IL-1, IL-6, and TGFβ1 playing a major role in driving the differentiation process,<sup>33</sup> while IL-22 is also sourced from Th2 cells, which are involved in immunity and remodeling.<sup>34</sup> Interleukin-17A, a member of the IL-17 cytokine family, is known to contribute to the host defense against pathogens in both mucosal and epithelial tissue; and IL-22, a member of the IL-10 cytokine family, plays a pivotal role in epithelial wound healing.<sup>35,36</sup>

Interleukin-17 and -22 are known to stimulate neutrophil infiltration and metalloproteinases (MMPs).<sup>37</sup> Studies have shown that IL-17 in particular contributes to disrupting the corneal barrier function due to the MMPs produced. This can have a large effect on the epithelial cell multilayer, causing erosion and surface irregularity.<sup>38</sup> Studies have also shown that IL-17RA (the receptor of IL-17A) is highly expressed in the






**FIGURE 6.** Expression of CK3 (A–H) and P63 (I–P) (shown in *green*) in cells on an in vitro corneal organ culture model under static (A–D, I–L) and rocking (E–H, M–P) conditions. (A, E, I, M) An unwounded cornea (control); (B, F, J, N) an unwounded cornea exposed to IL-17A; (C, G, K, O) a wounded cornea; (D, H, L, P) a wounded cornea exposed to IL-17A. Nuclei were counterstained with propidium iodide (*red*).

basal limbal region, suggesting that the region is susceptible to the inflammatory action of IL-17.<sup>39</sup> Donetti et al.<sup>40</sup> reported that IL-17 reduces proliferation, is destructive to tissue, and is described as proinflammatory.<sup>34</sup> Interleukin-22 has been reported to be responsible for the inhibition of differential gene expression while inducing proliferation and migration of keratinocytes.<sup>41</sup> Interleukin-17 alone has been implicated in dry eye,<sup>38</sup> scleritis, experimental autoimmune uveoretinitis,<sup>42</sup> herpetic stromal keratitis,<sup>43</sup> and multiple sclerosis,<sup>44</sup> while IL-17 and IL-22 together have been implicated in diseases related to skin inflammation, rheumatoid arthritis,<sup>44–46</sup> and uveitis.<sup>42</sup> This may explain the results in this study (Fig. 4) showing that IL-17 or the combination of IL-17 and -22 affected the epithelial morphology on the corneal surface.

In this particular study, LPS, which is found on the cell wall of Gram negative bacteria and is an endotoxin known to cause inflammatory responses to the host tissue,<sup>47,48</sup> had little or no effect on the appearance of the corneal model. Studies have shown that LPS may cause an inflammatory response only if

there is an entry site into the cornea, but certainly in these studies we failed to find any convincing response to LPS.

The major finding of this study was the extent to which subjecting corneas to gentle rocking improved the maintenance of the epithelium. Not only did they survive longer than under static conditions (4 weeks compared to 2 weeks), but there was a clear benefit when a wound was made in the model and it was allowed to recover. Recovery from wounding was very effective when corneas were kept under rocking conditions, much less so under static conditions. For the maturation of most 3D epithelial tissues in vitro, exposure to an air-liquid interface is essential, as is the case for skin, buccal mucosa, and esophageal mucosa.<sup>49–51</sup> However, in the case of the cornea, it has not been clear whether for in vitro experimentation it is best to keep the epithelia submerged or at an air-liquid interface or, as in this study, subject to intermittent bathing with media. This study clearly shows the benefit of subjecting these epithelia to the intermittent movement of media.



**FIGURE 7.** Expression of P63 in unwounded (A–C) and wounded (D–F) corneas cultured under rocking conditions. Images show P63 (green) expressed in nuclei (indicated by white arrows) in (A, D); nuclei counterstained with propidium iodide (red) in (B, E); and the merged images in (C, F).

When epithelia are injured, one of the first responses of the stromal cells is apoptosis of keratocytes, which can continue for 1 week after the injury has taken place.<sup>52</sup> It was noted in this study that most of the stromal cells close to the epithelial surface had died, giving a clear indication of the area where the wounding had taken place on the cornea during these experiments as shown in Figure 5. However, our attempts to confirm whether the stromal cells had undergone apoptosis were unsuccessful (results not shown), so we cannot make any conclusions regarding how these cells died.

With respect to expression of CK3 and P63, CK3 is known to be associated with the differentiated cells of the central corneal region,<sup>53</sup> while Pellegrini et al.<sup>54</sup> suggested that P63 is a stem cell marker expressed only in the basal layers of the limbal region. In characterization studies it was seen that wherever cells were present on the corneal epithelium, CK3 was expressed toward the central region of the cornea while

P63 was expressed at the periphery, which contains the limbus as well as the basal region of the corneal epithelium (Fig. 7).

We suggest that this relatively simple ex vivo rabbit cornea model can now more usefully be used to look at some of the influences on corneal repair and regeneration that are currently tested in live rabbit eyes.

### Acknowledgments

We thank the Wellcome Trust for funding this work through an Affordable Healthcare for India award. We thank Alison Weather from Hook Farm for supplying the rabbit eyes, Anthony Bullock, PhD, for advice on the use of inflammatory cytokines, and Frazer Bye, PhD, for his advice on the in-house rocking system.

Supported by the Wellcome Trust (Grant 091128).

Disclosure: **P. Deshpande**, None; **Í. Ortega**, None; **F. Sefat**, None; **V.S. Sangwan**, None; **N. Green**, None; **F. Claeysens**, None; **S. MacNeil**, None



## References

- Beebe DC. Maintaining transparency: a review of the developmental physiology and pathophysiology of two avascular tissues. *Semin Cell Dev Biol*. 2008;19:125-133.
- Kinoshita S, Adachi W, Sotozono C, et al. Characteristics of the human ocular surface epithelium. *Prog Retin Eye Res* 2001;20: 639-673.
- Fatt I, Bieber MT. The steady-state distribution of oxygen and carbon dioxide in the in vivo cornea: I. The open eye in air and the closed eye. *Exp Eye Res*. 1968;7:103-112.
- Ban Y, Cooper LJ, Fullwood NJ, et al. Comparison of ultrastructure, tight junction-related protein expression and barrier function of human corneal epithelial cells cultivated on amniotic membrane with and without air-lifting. *Exp Eye Res*. 2003;76:735-743.
- Satake Y, Dogru M, Yamane G, Kinoshita S, Tsubota K, Shimazaki J. Barrier function and cytologic features of the ocular surface epithelium after autologous cultivated oral mucosal epithelial transplantation. *Arch Ophthalmol*. 2008; 126:23-28.
- Dua HS, Gomes JA, Singh A. Corneal epithelial wound healing. *Br J Ophthalmol*. 1994;78:401-408.
- Li Z, Rumbaut RE, Burns AR, Smith CW. Platelet response to corneal abrasion is necessary for acute inflammation and efficient re-epithelialization. *Invest Ophthalmol Vis Sci*. 2006; 47:4794-4802.
- O'Brien TP, Li Q, Ashraf M, Matteson DM, Stark WJ, Chan C. Inflammatory response in the early stages of wound healing after excimer laser keratectomy. *Arch Ophthalmol*. 1998;116: 1470-1474.
- Li Z, Burns AR, Smith CW. Two waves of neutrophil emigration in response to corneal epithelial abrasion: distinct adhesion molecule requirements. *Invest Ophthalmol Vis Sci*. 2006;47: 1947-1955.
- Wilson SE, He Y-G, Weng J, et al. Epithelial injury induces keratocyte apoptosis: hypothesized role for the interleukin-1 system in the modulation of corneal tissue organization and wound healing. *Exp Eye Res*. 1996;62:325-338.
- Mohan RR, Kim WJ, Chen L, Wilson SE. Bone morphogenic proteins 2 and 4 and their receptors in the adult human cornea. *Invest Ophthalmol Vis Sci*. 1998;39:2626-2636.
- Gan L, Fagerholm P, Kim HJ. Effect of leukocytes on corneal cellular proliferation and wound healing. *Invest Ophthalmol Vis Sci*. 1999;40:575-581.
- Marrazzo G, Bellner L, Halilovic A, et al. The role of neutrophils in corneal wound healing in HO-2 null mice. *PLoS One*. 2011;6:e21180.
- Smith JA. Neutrophils, host defense, and inflammation: a double-edged sword. *J Leukoc Biol*. 1994;56:672-686.
- Deshpande P, Notara M, Bullett N, Daniels JT, Haddow DB, MacNeil S. Development of a surface-modified contact lens for the transfer of cultured limbal epithelial cells to the cornea for ocular surface diseases. *Tissue Eng Part A*. 2009;15:2889-2902.
- Deshpande P, Ramachandran C, Sefat F, et al. Simplifying corneal surface regeneration using a biodegradable synthetic membrane and limbal tissue explants. *Biomaterials*. 2013;34: 5088-5106.
- Notara M, Schrader S, Daniels J. The porcine limbal epithelial stem cell niche as a new model for the study of transplanted tissue-engineered human limbal epithelial cells. *Tissue Eng Part A*. 2011;17:741-750.
- Zhao B, Cooper LJ, Brahma A, MacNeil S, Rimmer S, Fullwood NJ. Development of a three-dimensional organ culture model for corneal wound healing and corneal transplantation. *Invest Ophthalmol Vis Sci*. 2006;47:2840-2846.
- Richard NR, Anderson JA, Weiss JL, Binder PS. Air/liquid corneal organ culture: a light microscopic study. *Curr Eye Res*. 1991;10:739-749.
- Foreman DM, Pancholi S, Jarvis-Evans J, McLeod D, Boulton MEA. Simple organ culture model for assessing the effects of growth factors on corneal re-epithelialization. *Exp Eye Res*. 1996;62:555-564.
- Cho K-A, Suh JW, Lee KH, Kang JL, Woo S-Y. IL-17 and IL-22 enhance skin inflammation by stimulating the secretion of IL-1 $\beta$  by keratinocytes via the ROS-NLRP3-caspase-1 pathway. *Int Immunol*. 2012;24:147-158.
- Andoh A, Zhang Z, Inatomi O, et al. Interleukin-22, a member of the IL-10 subfamily, induces inflammatory responses in colonic subepithelial myofibroblasts. *Gastroenterology*. 2005; 129:969-984.
- Cohn Z, Kim A, Huang L, Brand J, Wang H. Lipopolysaccharide-induced inflammation attenuates taste progenitor cell proliferation and shortens the life span of taste bud cells. *BMC Neurosci*. 2010;11:1-16.
- Eutamene H, Theodorou V, Schmidlin F, et al. LPS-induced lung inflammation is linked to increased epithelial permeability: role of MLCK. *Eur Respir J*. 2005;25:789-796.
- Berger RE, Corrsin S. A surface tension gradient mechanism for driving the pre-corneal tear film after a blink. *J Biomech*. 1974; 7:225-238.
- Schwab IR, Reyes M, Isseroff RR. Successful transplantation of bioengineered tissue replacements in patients with ocular surface disease. *Cornea*. 2000;19:421-426.
- Sangwan V, Matalia H, Vemuganti G, et al. Clinical outcome of autologous cultivated limbal epithelium transplantation. *Indian J Ophthalmol*. 2006;54:29-34.
- Ceccarini C, Eagle H. pH as a determinant of cellular growth and contact inhibition. *Proc Natl Acad Sci U S A*. 1971;68: 229-233.
- Potter SM, DeMarse TB. A new approach to neural cell culture for long-term studies. *J Neurosci Methods*. 2001;110:17-24.
- Zetterberg A, Engström W. Mitogenic effect of alkaline pH on quiescent, serum-starved cells. *Proc Natl Acad Sci U S A*. 1981; 78:4334-4338.
- Waite J, Skokos D. Th17 response and inflammatory autoimmune diseases. *Int J Inflam*. 2012;2012:819467.
- Damsker JM, Hansen AM, Caspi RR. Th1 and Th17 cells: adversaries and collaborators. *Ann N Y Acad Sci*. 2010;1183: 211-221.
- Chen Z, O'Shea J. Th17 cells: a new fate for differentiating helper T cells. *Immunol Res*. 2008;41:87-102.
- Eyerich S, Eyerich K, Pennino D, et al. Th22 cells represent a distinct human T cell subset involved in epidermal immunity and remodeling. *J Clin Invest*. 2009;119:3573-3585.
- Li Z, Burns AR, Miller SB, Smith CW. CCL20,  $\gamma\delta$  T cells, and IL-22 in corneal epithelial healing. *FASEB J*. 2011;25:2659-2668.
- Sabat R, Witte E, Witte K, Wolk K. IL-22 and IL-17: an overview. In: Quesniaux V, Ryffel B, Padova F, eds. *IL-17, IL-22 and Their Producing Cells: Role in Inflammation and Autoimmunity*. Basel: Springer; 2013:11-35.
- Stange J, Hepworth MR, Rausch S, et al. IL-22 mediates host defense against an intestinal intracellular parasite in the absence of IFN- $\gamma$  at the cost of Th17-driven immunopathology. *J Immunol*. 2012;188:2410-2418.
- De Paiva CS, Chotikavanich S, Pangelinan SB, et al. IL-17 disrupts corneal barrier following desiccating stress. *Mucosal Immunol*. 2009;2:243-253.
- Bian F, Qi H, Ma P, et al. An immunoprotective privilege of corneal epithelial stem cells against Th17 inflammatory stress by producing glial cell-derived neurotrophic factor. *Stem Cells*. 2010;28:2172-2181.

40. Donetti E, Cornaghi L, Gualerzi A, Baruffaldi Preis FW, Prignano F. An innovative three-dimensional model of normal human skin to study the proinflammatory psoriatic effects of tumor necrosis factor-alpha and interleukin-17. *Cytokine*. 2014;68:1-8.
41. Boniface K, Bernard F-X, Garcia M, Gurney AL, Lecron J-C, Morel F. IL-22 inhibits epidermal differentiation and induces proinflammatory gene expression and migration of human keratinocytes. *J Immunol*. 2005;174:3695-3702.
42. Amadi-Obi A, Yu C-R, Liu X, et al. TH17 cells contribute to uveitis and scleritis and are expanded by IL-2 and inhibited by IL-27/STAT1. *Nat Med*. 2007;13:711-718.
43. Maertzdorf J, Osterhaus AD, Verjans GM. IL-17 expression in human herpetic stromal keratitis: modulatory effects on chemokine production by corneal fibroblasts. *J Immunol*. 2002;169:5897-5903.
44. Jin W, Dong C. IL-17 cytokines in immunity and inflammation. *Emerg Microbes Infect*. 2013;2:e60.
45. Ikeuchi H, Kuroiwa T, Hiramatsu N, et al. Expression of interleukin-22 in rheumatoid arthritis: potential role as a proinflammatory cytokine. *Arthritis Rheum*. 2005;52:1037-1046.
46. Boniface K, Guignouard E, Pedretti N, et al. A role for T cell-derived interleukin 22 in psoriatic skin inflammation. *Clin Exp Immunol*. 2007;150:407-415.
47. Copeland S, Warren HS, Lowry SE, Calvano SE, Remick D. Inflammation and the host response to injury I. Acute inflammatory response to endotoxin in mice and humans. *Clin Diagn Lab Immunol*. 2005;12:60-67.
48. Schultz CL, Buret AG, Olson ME, Ceri H, Read RR, Morck DW. Lipopolysaccharide entry in the damaged cornea and specific uptake by polymorphonuclear neutrophils. *Infect Immun*. 2000;68:1731-1734.
49. Deshpande P, Ralston DR, MacNeil S. The use of allodermis prepared from Euro skin bank to prepare autologous tissue engineered skin for clinical use. *Burns*. 2013;39:1170-1177.
50. Bhargava S, Patterson JM, Inman RD, MacNeil S, Chapple CR. Tissue-engineered buccal mucosa urethroplasty—clinical outcomes. *Eur Urol*. 2008;53:1263-1271.
51. Green NH, Nicholls Z, Heath PR, et al. Pulsatile exposure to simulated reflux leads to changes in gene expression in a 3D model of oesophageal mucosa. *Int J Exp Pathol*. 2014;216-228.
52. Wilson SE, Mohan RR, Mohan RR, Ambrósio R Jr, Hong J, Lee J. The corneal wound healing response: cytokine-mediated interaction of the epithelium, stroma, and inflammatory cells. *Prog Retin Eye Res*. 2001;20:625-637.
53. Schermer A, Galvin S, Sun TT. Differentiation-related expression of a major 64K corneal keratin in vivo and in culture suggests limbal location of corneal epithelial stem cells. *J Cell Biol*. 1986;103:49-62.
54. Pellegrini G, Dellambra E, Golisano O, et al. p63 identifies keratinocyte stem cells. *Proc Natl Acad Sci U S A*. 2001;98:3156-3161.